The Gut Brain Axis - Exploring the Impact of Nutrition on Neurological Disorders

Sia Ittyavirah[1] Dr Soumya[2]
[1,2]Masters of Science in Food Nutrition and Dietetics
[1,2]Ms Ramaiah University of Applied Sciences, India

Abstract — In recent years, the interplay between the human gut and brain has emerged as a powerful paradigm in understanding health and disease. The gut-brain axis (GBA), is a communication network linking the central nervous system with the gastrointestinal system, which is no longer viewed solely through the lens of digestion or neurotransmission; it is now considered fundamental to emotional regulation, immune balance, and even cognitive performance. One of the most exciting frontiers within this field lies in the influence of the gut microbiota, the trillions of microorganisms residing in our digestive tract that interact closely with our nervous and immune systems.[1], [2]. As modern science delves into the roles of these microbes, a consistent theme has emerged that diet is one of the most significant and immediate modulators of gut microbial health. The food we consume can rapidly alter the structure and function of our microbiota, leading to cascading effects on brain health. For instance, fiber-rich diets promote the production of short-chain fatty acids (SCFAs), which have anti-inflammatory and neuroprotective properties. Meanwhile, processed and high-fat diets can disrupt microbial balance, leading to inflammation, impaired gut barrier function, and altered neurotransmitter signaling [3]. Neurological disorders such as depression, anxiety, Attention-Deficit/Hyperactivity Disorder (ADHD), autism spectrum disorder, and even neurodegenerative diseases like Alzheimer's and Parkinson's have all shown links to gut dysbiosis. While traditional treatments focus on pharmacological approaches, the potential to influence brain function through dietary change offers a promising, non-invasive avenue for both prevention and adjunctive care. Moreover, microbiota directed therapies such as probiotics, prebiotics, and ketogenic diets are gaining attention for their ability to support neurological outcomes through gut modulation [4]. This review aims to explore how various dietary patterns, nutrients, and microbiota-based interventions influence the gut-brain axis and impact the onset or progression of neurological disorders. By compiling current findings, the paper seeks to identify both well-supported mechanisms and key areas that demand further investigation, ultimately contributing to a more integrated understanding of brain health through the lens of nutrition and gut microbiology [5].

Keywords: Gut-Brain Axis (GBA), Microbiota-Gut-Brain Communication, Gut Microbiota, Diet and Brain Health, Short-Chain Fatty Acids (SCFAs), Neuroinflammation, Neurotransmitter Signaling, Gut Dysbiosis

I. Introduction

The gut-brain axis (GBA) is a complex, bidirectional communication system that connects the central nervous system (CNS) and the enteric nervous system (ENS), allowing the brain and the gastrointestinal tract to influence each other's function. This axis operates through neural, endocrine, immune, and metabolic signaling pathways, and is increasingly recognized as a critical regulator of neurodevelopment, behavior, and emotional well-being [6].

Central to the GBA is the gut microbiota, a dynamic community of microorganisms residing in the gastrointestinal tract. These microbes perform essential roles in digestion, immune modulation, and the synthesis of neuroactive compounds such as serotonin, gamma-aminobutyric acid (GABA), and short-chain fatty acids (SCFAs), all of which can influence brain function. Emerging evidence shows that alterations in gut microbiota composition—often referred to as dysbiosis—can disrupt this axis, contributing to the development and progression of various neurological and psychiatric disorders, including depression, anxiety, ADHD, autism spectrum disorder (ASD), and neurodegenerative diseases like Alzheimer's and Parkinson's [6].

Among the most powerful and modifiable factors influencing the gut microbiome is diet. Nutritional intake directly shapes the composition and function of gut microbiota, with immediate and long-term consequences for host health. While Western diets, high in saturated fats, processed foods, and added sugars, are associated with reduced microbial diversity and inflammation, other dietary patterns such as the Mediterranean and ketogenic diets promote microbial environments more conducive to brain health [7].

The purpose of this review is to critically examine how dietary patterns, specific nutrients, and microbiota-targeted interventions such as probiotics, prebiotics, fermented foods, and polyphenols influence the GBA and contribute to the management or prevention of neurological disorders. By exploring the mechanisms linking nutrition, the microbiota, and brain function, this review aims to highlight the potential of diet-based strategies in promoting neurological resilience and guiding future therapeutic approaches. [5].

II. Mechanism of Gut - Brain Axis

"The gut-brain axis is an intricate network of signaling pathways that facilitate constant communication between the gastrointestinal tract and the central nervous system. This axis integrates neural, immune, endocrine, and metabolic signaling pathways, enabling the brain to influence gastrointestinal function while also allowing the gut to impact mood, cognition, and behavior. Far from being an isolated digestive organ, the gut is now recognized as a key contributor in neurophysiological health. This recognition has prompted in-depth exploration of the multiple communication mechanisms that constitute the gut-brain axis [4].

One of the most direct and well-studied pathways in this communication system is the vagus nerve. As the longest

cranial nerve in the body, it serves as a primary neural channel for facilitating bidirectional communication between the gut and the brain.Sensory neurons in the gut wall detect mechanical stretching, microbial metabolites, and chemical stimuli. These signals are then transmitted to brain regions such as the brainstem and limbic system.The vagus nerve also carries efferent signals that regulate processes like gut motility, secretion, and inflammatory responses, forming a feedback loop between the two systems [8].

In parallel with this neural communication, the immune system plays a vital role in relaying information from the gut to the brain. The intestinal lining contains a diverse array of immune cells that constantly monitor the gut environment for pathogens and microbial disturbances. When imbalances or infections occur, these cells release cytokines, small signaling proteins that influence inflammation and can travel through the bloodstream to reach the brain. Elevated levels of pro-inflammatory cytokines have been associated with changes in mood, cognition, and neurodegeneration. Chronic low-grade inflammation originating in the gut has been proposed as a contributing factor in psychiatric disorders such as depression, anxiety, and even neurodegenerative diseases.Thus, immune activity in the gut not only defends against pathogens but also plays a pivotal role in modulating brain function and emotional states [9].

A third essential pathway is hormonal signaling, particularly through the hypothalamic-pituitary-adrenal (HPA) axis, the body's central stress response system. Psychological or physical stressors activate the HPA axis, leading to the release of cortisol, a glucocorticoid hormone. Cortisol influences gut barrier integrity, alters the composition of the gut microbiota, and modulates immune responses. Over time, stress-induced changes in the gut environment can feedback into the brain, potentially impairing emotional regulation and stress resilience [10].

Crucially, these systems are influenced and modulated by the gut microbiota—a diverse and densely populated microbial community residing in the colon. These microbes have a profound influence on neurodevelopment, synaptic plasticity, and behavioral responses. During early life, the composition and diversity of the microbiota contribute to critical windows of brain development, affecting processes such as myelination, neurotransmitter receptor expression, and neural circuit formation. Disruption of this early microbial colonization has been associated with long-term consequences for mental health and stress regulation [11].

A key way in which microbes exert their influence on the brain is through the production of bioactive metabolites. Among the most important are short-chain fatty acids (SCFAs) primarily acetate, propionate, and butyrate produced by bacterial fermentation of dietary fiber. These SCFAs can cross the blood-brain barrier and are known to regulate gene expression, reduce neuroinflammation, support the structural integrity, and can also influence brain-derived neurotrophic factor (BDNF) levels, thereby supporting synaptic plasticity and neuroprotection [12]. Butyrate, in particular, has been studied for its neuroprotective and anti-inflammatory properties. Also certain bacteria within the gut are capable of synthesizing or modulating neurotransmitters that are also active in the human brain. For example, a large portion of serotonin is produced in the gut, where it regulates intestinal motility, but its precursors can influence central serotonin levels. Other microbial species produce or affect levels of gamma-aminobutyric acid (GABA) , the brain's main inhibitory neurotransmitter, as well as dopamine and acetylcholine, which are involved in motivation, learning, and memory [13].

Collectively, these interconnected pathways highlight the profound bidirectional influence between gut and brain. Disruptions whether from poor diet, stress, infection, or microbial imbalance can cascade through these systems, affecting mental health. Conversely, interventions targeting gut health offer promising avenues for enhancing neurological and psychological well-being.



Fig. 1: Diagram showing bidirectional communication between gut and brain through vagus nerve, immune, endocrine, and metabolic pathways

III. Role of Nutrition in Shaping Gut Microbiome

The gut microbiome comprises trillions of microorganisms, including bacteria, archaea, viruses, and fungi, residing in the human gastrointestinal tract. This dynamic and symbiotic ecosystem plays a crucial role in digestion, vitamin synthesis, immune function, and neurophysiological regulation. A diverse and balanced microbiota is associated with optimal health, whereas disruptions in this balance are increasingly linked to a wide range of chronic diseases including obesity, irritable bowel syndrome (IBS), metabolic syndrome, and neuropsychiatric disorders such as anxiety and depression [14].

Among the many external factors influencing gut health, diet stands out as one of the most powerful modulators of the microbiome. Dietary components can rapidly and profoundly alter microbial composition and function often within days underscoring the microbiome's responsiveness to nutritional inputs [15]. The concept of eubiosis refers to a state of balanced, diverse, and health-promoting gut microbiota, while dysbiosis describes a state marked by reduced diversity, dominance of pathogenic species, and impaired metabolic function. Diet-induced dysbiosis is now recognized as a key contributor to chronic inflammation, metabolic disruption, and altered gut-brain communication.

A. Dietary Patterns, Microbiome Composition: Western Diet vs. Mediterranean Diet

The Western dietary pattern, characterized by high intake of refined carbohydrates, saturated fats, red and processed meats, added sugars, and low dietary fiber, is associated with a reduction in microbial diversity and an increase in opportunistic and pro-inflammatory species such as Bacteroides and Firmicutes. This dietary pattern fosters dysbiosis and contributes to low-grade systemic inflammation, metabolic endotoxemia, and neuroinflammation all of which may underlie various metabolic and neurobehavioral disorder [16].

In contrast, the Mediterranean diet, which is rich in fruits, vegetables, legumes, whole grains, nuts, olive oil, polyphenols, and fermented foods, supports microbial diversity and promotes the abundance of beneficial taxa, including Bifidobacteria, Lactobacillus, and Faecalibacterium prausnitzii [7].These microbes are associated with increased production of short-chain fatty acids (SCFAs) like butyrate, which exert anti-inflammatory, gut-barrier-strengthening, and neuroprotective effects. Notably, interventional studies have demonstrated that shifting from a Western to a Mediterranean diet can induce measurable microbiota changes in as little as 3–days, with increased SCFA production and improved metabolic parameters [17].

B. Dietary Patterns, Microbiome Composition: Ketogenic Diet

The ketogenic diet (KD), a high-fat, low-carbohydrate, and moderate-protein dietary regimen, has gained prominence not only as a metabolic intervention for obesity and type diabetes, but also for its potential in neurological and inflammatory disorders. Central to its mechanism of action is a shift in energy metabolism toward ketone body production (notably β-hydroxybutyrate), which has been shown to exert anti-inflammatory, neuroprotective, and mitochondrial-enhancing effects. More recently, studies have highlighted the gut microbiome as a key mediator of the KD's therapeutic effects.

Research has shown that the ketogenic diet (KD) causes clear changes in the gut microbiome. Studies conducted on mouse show KD to increase the levels of beneficial bacteria like *Akkermansia muciniphila* and *Parabacteroides*, while reducing carbohydrate-fermenting species such as *Bifidobacterium* and *Lactobacillus*. These microbial changes were found to be essential for the diet's anti-seizure effects, as mice given antibiotics or raised in germ-free environments did not experience seizure protection despite having elevated ketone levels [18]. Likewise, a KD enriched with medium-chain triglycerides (MCTs) was shown to improve gut barrier function and reduce inflammation. This was associated with an increase in *Akkermansia* species, which help maintain the intestinal mucus layer and strengthen tight junctions between gut cells both of which are important for preventing "leaky gut" and limiting the movement of inflammatory toxins like lipopolysaccharides [19].

However, strict ketogenic diets often lead to a decrease in microbial diversity (alpha diversity), likely due to low fiber intake. The effects of this reduced diversity are complex and may vary based on the types of fats consumed, the inclusion of non-digestible fibers, and the individual's baseline microbiota composition.

Ketogenic diets promote the production of ketone bodies such as β-hydroxybutyrate (BHB), which have been shown to inhibit histone deacetylases, reduce oxidative stress, and modulate inflammatory pathways. BHB also influences the gut microbiome by altering the intestinal luminal environment, favoring microbial taxa adapted to a high-fat, low-carbohydrate milieu [19].

Effect	Associated microbial changes	Implication
↓Carbohydrate intake	↓ Bifidobacterium, Lactobacillus	Reduced fermentation; lower SCFA
↑ Fat intake (esp. MCTs)	↑ Akkermansia muciniphila, Parabacteroides	Improved barrier, anti-inflammatory
↓ Fiber intake	↓ SCFA (butyrate)	Weaker epithelial support (may be mitigated)
↓ α-diversity	Varies	May affect long-term resilience
↑ Ketones	Indirect microbial modulation	Improved neurochemical signaling

Table 1: Microbiome Changes and Effect

C. Specific Dietary Components and Their Impact on the Microbiome

1) Dietary Fiber

Fiber is abundant in plant-based foods and serves as the primary fuel for gut bacteria that produce SCFAs such as acetate, propionate, and butyrate. These metabolites help maintain gut integrity, reduce inflammation, and interact with the central nervous system. A fiber-rich diet enhances the abundance of beneficial taxa like *Bifidobacteria* and *Faecalibacterium prausnitzii*, which are commonly depleted in individuals with depression and neurodegenerative diseases. As a prebiotic, fiber selectively stimulates the growth of beneficial gut bacteria like *Bifidobacterium* and *Lactobacillus*. The fermentation of fiber in the colon yields SCFAs, which play a central role in gut integrity, anti-inflammatory signaling, and metabolic health [20].

2) Polyphenols

Polyphenols are bioactive compounds found in plant-based foods like berries, tea, cocoa, and red wine, which also possess prebiotic properties. They modulate the gut microbiota by inhibiting pathogenic bacteria and promoting beneficial species such as *Akkermansia muciniphila*. Polyphenols are metabolized by gut bacteria into bioactive compounds that influence oxidative stress, neuronal signaling, and inflammation. Regular intake of polyphenol-rich foods can selectively support the growth of bacteria that modulate the immune system and produce neuroprotective agents [21].

3) Fermented Foods

Fermented foods such as yogurt, kefir, kimchi, tempeh and miso introduce live microbial cultures (probiotics) into the gut. Regular consumption of fermented foods has been shown to increase microbial diversity, lower gut pH, and enhance the

gut-brain axis, potentially reducing symptoms of anxiety and depression [22].

4) Sugar and Artificial Sweeteners

High intake of sugar diets can encourage the growth of pathogenic bacteria and fungi, increase gut permeability (leaky gut), and trigger systemic inflammation.These dietary components may reduce populations of beneficial species and disrupt microbial signaling pathways involved in brain health and immune modulation, promoting the growth of inflammatory and pathogenic microbes. Additionally, non-nutritive sweeteners such as sucralose and aspartame may disrupt the microbiome and impair glucose metabolism [23].

Fats

5) Dietary fats differentially affect the microbiome:

Saturated and trans fats: Linked to increased levels of endotoxin-producing bacteria and reduced microbial diversity. Excessive fat intake, particularly from animal sources, has also been linked to a reduction in *Bacteroidetes* and an increase in *Firmicutes*, a microbial shift commonly associated with obesity and metabolic dysfunction.

Unsaturated fats, especially omega-3 fatty acids, eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), found in fatty fishlike salmon and mackerel, exert anti-inflammatory and neuroprotective effects. These fats can modulate gut microbial composition, promoting the growth of SCFA-producing bacteria and improving the integrity of the intestinal barrier. Additionally, omega-3s have been shown to improve symptoms of depression, anxiety, and cognitive decline through their action on neuronal membranes and inflammatory pathways [24].

D. How Diet Promotes Eubiosis or Dysbiosis

1) Eubiosis

- SCFA Production: SCFAs, especially butyrate, maintain epithelial integrity and suppress pro-inflammatory cytokines.
- Barrier Function: Nutrient-driven upregulation of tight junction proteins strengthens the gut barrier.
- Immune Modulation: Balanced microbiota promote tolerogenic immune responses.
- Neurochemical Signaling: Diet influences microbial production of neurotransmitters (e.g., serotonin, dopamine, GABA).

2) Dysbiosis mechanism

- Loss of Microbial Diversity: Reduced ability to resist pathogens or metabolize nutrients effectively.
- Leaky Gut: Increased gut permeability allows endotoxins to enter systemic circulation.
- Systemic Inflammation: Chronic activation of inflammatory pathways.
- Gut-Brain Axis Disruption: Altered microbiota impair neurotransmitter synthesis and stress response systems.

IV. GUT - BRAIN AXIS AND SPECIFIC NEUROLOGICAL DISORDERS

A growing body of evidence supports the hypothesis that disturbances in the gut microbiota can influence the onset and progression of several neurological disorders via the gut-brain axis. Alterations in microbial diversity, metabolite production, and gut permeability have all been linked to neuroinflammatory and neurobehavioral changes observed in various mental and neurodevelopmental conditions. By modulating immune responses, neurotransmitter synthesis, and stress pathways, the gut microbiota emerges as a potential biomarker and therapeutic target for a range of disorders affecting emotional regulation, attention, and cognition [25].

A. Stress, Depression and Anxiety

1) Evidence Linking Dysbiosis with Mood Disorders

Individuals with major depressive disorder (MDD) or generalized anxiety disorder (GAD) exhibit distinct microbial profiles compared to healthy individuals. These changes often involve reduced microbial diversity, along with lower levels of beneficial genera and an increase in pro-inflammatory taxa, including certain *Proteobacteria* [26].

One of the key mechanisms linking gut dysbiosis to mood disorders is increased intestinal permeability. This condition allows endotoxins such as lipopolysaccharide (LPS) to translocate from the gut lumen into systemic circulation, triggering low-grade systemic inflammation and neuroinflammatory responses. Elevated levels of circulating pro-inflammatory cytokines—including interleukin-6 (IL-6), tumor necrosis factor alpha (TNF-α), and C-Reactive Protein (CRP) have been consistently associated with depression and anxiety in both clinical and animal studies [27].

Experimental research using germ-free (GF) animal models, which lack an indigenous gut microbiota, has shown the role of the microbiota in modulating stress physiology. These animals display exaggerated hypothalamic-pituitary-adrenal (HPA) axis activation, elevated corticosterone levels, and heightened anxiety-like behaviors. Colonization with specific bacterial strains can reverse these abnormalities, suggesting a causal link between gut microbiota and emotional regulation [28].

B. Attention-Deficit/Hyperactivity Disorder

1) Gut Microbiota Alterations in ADHD

Attention-Deficit/Hyperactivity Disorder is a neurodevelopmental condition characterized by symptoms of inattention, impulsivity, and hyperactivity. Although its etiology is multifactorial—encompassing genetic, environmental, and neurobiological factors—emerging research points to a potential role for the gut-brain axis in modulating neurodevelopmental outcomes, including those seen in ADHD [29].

Dietary factors are believed to have a significant impact on ADHD symptoms, both through direct nutritional effects on brain function and through modulation of the gut microbiota. Diets high in refined sugars, artificial additives, and low in fiber have been associated with a higher prevalence of ADHD. Conversely, diets rich in omega-3 fatty acids, polyphenols, and plant fibers may help support microbial diversity and reduce behavioral symptoms [29].

Probiotic and prebiotic supplementation has been explored in the context of ADHD, with some promising but limited evidence. One randomized controlled trial found that early-life administration of *Lactobacillus rhamnosus* reduced the risk of developing ADHD and Asperger's by age 13. Other studies suggest that multispecies probiotics may help improve attention and reduce impulsivity [30].

C. Alzheimer's Disease and Cognitive Decline

1) Role of Inflammation and Oxidative Stress

The gut microbiota has a crucial role in the development and progression of Alzheimer's disease (AD), particularly through its influence on inflammation and oxidative stress—two key pathological mechanisms underlying neurodegeneration. The term gut dysbiosis which has been increasingly associated with systemic and cerebral pathological changes seen in AD [31].

One primary mechanism by which dysbiosis contributes to AD is the promotion of systemic inflammation. Certain gut microbial profiles, characterized by decreased levels of beneficial anti-inflammatory bacteria (e.g., *Bifidobacterium* and *Faecalibacterium*) and increased levels of pro-inflammatory taxa (e.g., *Escherichia/Shigella*), have been found in AD patients [32]. These pro-inflammatory bacteria can stimulate the release of cytokines such as IL-1β, IL-6, and TNF-α, which not only contribute to systemic inflammation but may also penetrate or compromise the blood-brain barrier (BBB). A disrupted BBB allows these peripheral immune molecules and microbial products, such as lipopolysaccharides (LPS) and amyloid-like bacterial proteins, to access the brain parenchyma, where they promote microglial activation, neuroinflammation, and β-amyloid deposition [33].

Another key process influenced by gut dysbiosis is oxidative stress, a condition defined by an imbalance between the production of reactive oxygen species (ROS) and the body's antioxidant defenses. In the context of AD, oxidative stress contributes to lipid peroxidation, protein misfolding, and DNA damage, all of which are implicated in the deterioration of neuronal integrity and function. Altered gut microbiota may elevate oxidative stress by reducing levels of antioxidant-producing bacteria or through the increased production of ROS-inducing metabolites (Bonfili et al., 2020). Moreover, the oxidative damage resulting from microbial dysregulation may further exacerbate the pathological accumulation of tau protein tangles and β-amyloid plaques, two defining histopathological hallmarks of Alzheimer's disease [34].

Recent preclinical models suggest that modulating the gut microbiota—via antibiotics, probiotics, or fecal microbiota transplantation—can attenuate both inflammation and oxidative damage in the brain and improve cognitive outcomes, reinforcing the causal relationship between the gut and neurodegenerative processes. These findings support the hypothesis that gut dysbiosis serves as a chronic trigger of both peripheral and central immune responses and oxidative injury, thereby accelerating neurodegeneration and cognitive decline in Alzheimer's disease [35].

D. Autism Spectrum Disorder (ASD)

1) Gut Dysbiosis in Children with ASD

Children diagnosed with autism spectrum disorder (ASD) frequently exhibit gastrointestinal (GI) abnormalities, including constipation, diarrhea, abdominal pain, and bloating. These symptoms are not only common but are often correlated with the severity of core behavioral symptoms such as irritability, repetitive behaviors, and social withdrawal. Emerging evidence suggests that these GI symptoms may be rooted in alterations of the gut microbiota, also referred to as gut dysbiosis [36].

Studies utilizing 16S rRNA sequencing and metagenomics have consistently reported increased abundance of potentially pathogenic bacterial species, particularly Clostridium spp., in children with ASD, alongside decreased levels of beneficial microbes, such as *Bifidobacterium, Lactobacillus,* and *Prevotella*. These alterations disrupt intestinal barrier integrity and facilitate the translocation of bacterial endotoxins into systemic circulation, triggering immune activation and low-grade systemic inflammation, which may extend to the brain and influence neurodevelopmental processes [37].

Increased intestinal permeability has been proposed as a central mechanism linking microbial imbalance to neurological dysfunction in ASD. Dysbiotic gut communities may alter the production of short-chain fatty acids (SCFAs), such as butyrate and propionate, which have been known for its neuromodulatory effects. Elevated levels of propionic acid, in particular, have been associated with abnormal behavior and mitochondrial dysfunction in rodent models that mimic features of autism [38].

E. Parkinson's Disease

1) Constipation and Microbiota as Early Indicators

Parkinson's disease (PD) is a progressive neurodegenerative disorder primarily characterized by motor symptoms such as tremors, bradykinesia, and rigidity. However, non-motor symptoms, particularly gastrointestinal disturbances like constipation, often precede motor manifestations by several years. This has led to growing interest in the gut-brain axis and the role of gut microbiota in the early pathogenesis of PD [39].

Chronic constipation, sometimes appearing decades before PD diagnosis, is now recognized as a potential prodromal marker of the disease. Studies have revealed significant alterations in the gut microbial communities of individuals with PD. These alterations include reduced abundance of SCFA-producing bacteria such as *Faecalibacterium prausnitzii* and *Roseburia*, along with an increase in pro-inflammatory microbes, including certain *Enterobacteriaceae*. The reduction in butyrate, a short-chain fatty acid (SCFA) with anti-inflammatory and neuroprotective properties, is thought to impair the intestinal barrier and promote systemic and neuroinflammation [40].

Studies have shown that restoring SCFA levels through dietary fiber, prebiotics, or direct supplementation with butyrate can better motor symptoms and reduce neuroinflammatory markers in animal models of PD. These findings suggest a potential therapeutic role for targeting gut microbiota to modulate disease progression [41].

V. Conclusion

The collective evidence reviewed in this article underscores the pivotal role of the gut microbiota in shaping neurological health through the gut-brain axis. Far from being isolated systems, the gut and brain communicate continuously via a range of biological signals—many of which are influenced by dietary choices and microbial metabolites. As such, the gut microbiota represents both a reflection of long-term dietary

patterns and a mechanistic conduit through which nutrition can influence cognitive and emotional outcomes.

Eubiosis, characterized by microbial diversity and functional richness, supports a well-regulated immune system, robust intestinal barrier, and balanced neurochemical signaling. In contrast, dysbiosis, often triggered by poor dietary habits, can lead to systemic inflammation, altered neurotransmitter levels, and increased vulnerability to neurological and psychiatric conditions. Nutritional strategies that restore microbial balance—such as increasing fiber and polyphenol intake, incorporating fermented foods and probiotics, and limiting processed sugars and unhealthy fats—show promise in reducing symptoms of anxiety, depression, ADHD, and possibly slowing the progression of neurodegenerative diseases.

Despite these advances, several research gaps remain. More longitudinal, human-based studies are needed to establish causal relationships between specific diets, microbial changes, and mental health outcomes. There is also a need to personalize dietary recommendations based on individual microbiome profiles, genetic backgrounds, and lifestyle factors. Additionally, the therapeutic potential of next-generation probiotics, postbiotics, and microbiome-targeted interventions (such as fecal microbiota transplantation) warrants deeper exploration.

In addition, future research should explore the role of early-life nutrition in shaping long-term brain health, especially during critical developmental windows. Digital health tools like wearable biosensors and AI-powered diet tracking apps may revolutionize personalized gut-brain axis monitoring and interventions. Furthermore, understanding the complex interactions between microbiota, the immune system, and neuroendocrine signaling can unlock novel therapeutic targets. Investigating the effects of commonly prescribed medications on microbiota composition may also reveal strategies to enhance the efficacy of dietary and psychobiotic therapies.

In conclusion, the gut-brain axis offers a powerful framework for rethinking the role of nutrition in mental and neurological health. By modulating the gut microbiota through strategic dietary interventions, it may be possible to improve brain function, enhance emotional well-being, and prevent or alleviate symptoms of neurological disorders. Future research and clinical practice would benefit from embracing this integrative, microbiome-informed approach to nutrition and neuroscience.

REFERENCES

[1] G. Sharon, T.R. Sampson, D.H. Geschwind, and S.K. Mazmanian, "The Central Nervous System and the Gut Microbiome," Cell, vol. 167, pp. 915–932, Nov. 2016.
[2] E.A. Mayer, R. Knight, S.K. Mazmanian, J.F. Cryan, and K. Tillisch, "Gut microbes and the brain: paradigm shift in neuroscience," Journal of Neuroscience, vol. 35, pp. 13864–13871, Oct. 2015.
[3] K. Martinez-Guryn et al., "Small intestine microbiota regulate host digestive and absorptive adaptive responses to dietary lipids," Cell Host & Microbe, vol. 23, pp. 458–469, Apr. 2018.
[4] J.F. Cryan, K.J. O'Riordan, K. Sandhu, V. Peterson, and T.G. Dinan, "The microbiota-gut-brain axis," Physiological Reviews, vol. 99, pp. 1877–2013, Oct. 2019.
[5] T.G. Dinan and J.F. Cryan, "Gut-brain axis in 2016: brain-gut-microbiota axis—mood, metabolism and behaviour," Nature Reviews Gastroenterology & Hepatology, vol. 14, pp. 69–70, Feb. 2017.
[6] M.H. Mohajeri, G. La Fata, R.E. Steinert, and P. Weber, "Relationship between the gut microbiome and brain function," Nutrition Reviews, vol. 76, pp. 481–496, Jul. 2018.
[7] F. De Filippis et al., "High-level adherence to a Mediterranean diet beneficially impacts the gut microbiota and associated metabolome," Gut, vol. 65, pp. 1812–1821, Nov. 2016.
[8] B. Bonaz, T. Bazin, and S. Pellissier, "The vagus nerve at the interface of the microbiota–gut–brain axis," Frontiers in Neuroscience, vol. 12, pp. 1–17, Apr. 2018.
[9] A.M. De Miranda, T. Milenkovic, S.R. Kennedy, et al., "The gut–immune–brain axis: Functions and clinical implications," Current Opinion in Pharmacology, vol. 48, pp. 23–29, Apr. 2018.
[10] J.A. Foster, L. Rinaman, and J.F. Cryan, "Stress and the gut-brain axis: regulation by the microbiome," Neurobiology of Stress, vol. 7, pp. 124–136, Jan. 2017.
[11] P. Luczynski et al., "Adult microbiota-deficient mice have distinct dendritic morphological changes," Brain, Behavior, and Immunity, vol. 68, pp. 179–193, Mar. 2016.
[12] B. Dalile, L. Van Oudenhove, B. Vervliet, and K. Verbeke, "The role of short-chain fatty acids in microbiota–gut–brain communication," Nature Reviews Gastroenterology & Hepatology, vol. 16, pp. 461–478, Aug. 2019.
[13] P. Strandwitz et al., "GABA-modulating bacteria of the human gut microbiota," Nature Microbiology, vol. 4, pp. 396–403, Mar. 2018.
[14] A.B. Shreiner, J.Y. Kao, and V.B. Young, "The gut microbiome in health and in disease," Current Opinion in Gastroenterology, vol. 31, pp. 69–75, Jan. 2015.
[15] L.A. David et al., "Diet rapidly and reproducibly alters the human gut microbiome," Nature, vol. 505, pp. 559–563, Jan. 2014.
[16] M.K. Zinöcker and I.A. Lindseth, "The Western diet–microbiome-host interaction and its role in metabolic disease," Nutrients, vol. 10, no. 3, pp. 365, Mar. 2018.
[17] C. Zhu et al., "Human gut microbiome responds to short-term interventions," Molecular Nutrition & Food Research, vol. 64, pp. e1900672, Jan. 2020.
[18] C.A. Olson et al., "The gut microbiota mediates the anti-seizure effects of the ketogenic diet," Cell, vol. 173, pp. 1728–1741, Jun. 2018.
[19] Q.Y. Ang et al., "Ketogenic diets alter the gut microbiome resulting in decreased intestinal Th17 cells," Cell Metabolism, vol. 32, pp. 816–833, Nov. 2020.
[20] A. Koh, F. De Vadder, P. Kovatcheva-Datchary, and F. Bäckhed, "Role of short-chain fatty acids in host physiology," Cell, vol. 165, pp. 1332–1345, Jun. 2016.
[21] D.E. Roopchand et al., "Dietary polyphenols promote growth of beneficial gut microbiota," Journal of

Nutritional Biochemistry, vol. 26, pp. 1425–1432, Dec. 2015.
[22] M.E. Sanders, D.J. Merenstein, C.A. Merrifield, and R. Hutkins, "Probiotics for human use," Gut Microbes, vol. 14, no. 1, pp. 1–18, 2023.
[23] J. Suez et al., "Artificial sweeteners induce glucose intolerance by altering the gut microbiota," Nature, vol. 514, pp. 181–186, Oct. 2014.
[24] C.A. Simpson et al., "The gut microbiota in anxiety and depression – A review," Current Opinion in Psychiatry, vol. 34, pp. 611–617, Sep. 2021.
[25] T.G. Dinan, C. Stanton, and J.F. Cryan, "Psychobiotics: A novel class of psychotropic," Biological Psychiatry, vol. 74, pp. 720–726, Oct. 2013.
[26] J.R. Kelly et al., "Transferring the blues: Depression-associated gut microbiota induces neurobehavioural changes," Molecular Psychiatry, vol. 21, pp. 1281–1288, Aug. 2016.
[27] M. Maes, M. Kubera, and J.C. Leunis, "The gut-brain barrier in major depression," Neuro Endocrinology Letters, vol. 33, no. 1, pp. 103–110, 2012.
[28] K.M. Neufeld, N. Kang, J. Bienenstock, and J.A. Foster, "Effects of intestinal microbiota on anxiety-like behavior," Neurogastroenterology & Motility, vol. 23, no. 3, pp. 255–e119, Mar. 2011.
[29] J. Szopinska-Tokov et al., "Gut microbiota in ADHD: Relevance and therapeutic targets," Microbiome, vol. 11, no. 1, pp. 59, Apr. 2023.
[30] A. Pärtty, M. Kalliomäki, P. Wacklin, S. Salminen, and E. Isolauri, "A possible link between early probiotic intervention and the risk of neuropsychiatric disorders," Pediatric Research, vol. 77, pp. 823–828, Jun. 2015.
[31] P. Liu et al., "Altered microbiomes distinguish Alzheimer's disease from amnestic mild cognitive impairment," Journal of Translational Medicine, vol. 18, no. 1, pp. 30, Jan. 2020.
[32] A. Cattaneo et al., "Association of brain amyloidosis with pro-inflammatory gut bacterial taxa and peripheral inflammation markers in cognitively impaired elderly," Journal of Alzheimer's Disease, vol. 58, no. 1, pp. 1–14, 2017.
[33] Z.Q. Zhuang et al., "Gut microbiota is altered in patients with Alzheimer's disease," Journal of Alzheimer's Disease, vol. 63, no. 4, pp. 1337–1346, 2018.
[34] L. Bonfili et al., "Microbiota modulation counteracts Alzheimer's disease progression influencing neuronal proteolysis and gut hormones plasma levels," FASEB Journal, vol. 34, no. 1, pp. 645–661, Jan. 2020.
[35] T. Harach et al., "Reduction of Abeta amyloid pathology in APPPS1 transgenic mice in the absence of gut microbiota," Scientific Reports, vol. 7, pp. 41802, Feb. 2017.
[36] C. Holingue et al., "Gastrointestinal symptoms in autism spectrum disorder: A review of the literature on ascertainment and prevalence," Autism Research, vol. 11, pp. 24–36, Jan. 2018.
[37] D.W. Kang et al., "Microbiota transfer therapy alters gut ecosystem and improves gastrointestinal and autism symptoms: an open-label study," Microbiome, vol. 5, pp. 10, Jan. 2017.
[38] D.F. MacFabe, "Short-chain fatty acid fermentation products of the gut microbiome: implications in autism spectrum disorders," Microbial Ecology in Health and Disease, vol. 27, pp. 1–13, Dec. 2016.
[39] M.G. Cersosimo et al., "Gastrointestinal manifestations in Parkinson's disease: prevalence and pathophysiology," Movement Disorders Clinical Practice, vol. 9, no. 1, pp. 54–62, Jan. 2022.
[40] J.R. Bedarf et al., "Functional implications of microbial and viral gut metagenome changes in early-stage Parkinson's disease," Genome Medicine, vol. 9, pp. 39, Apr. 2017.
[41] S. Sharma et al., "Biomarkers in Parkinson's disease," Translational Neurodegeneration, vol. 4, pp. 1–11, 2015.

