

During 5th Instar Development a Correlative Study on Changes in Concentration of Haemolymph Ascorbic Acid of Bivoltine Races of *Bombyx Mori.L*

Dr. Amresh Pattnaik¹ Prof. Subasini Pattnaik²

¹Research Scholar ²Professor

^{1,2}Department of Zoology

^{1,2}Berhampur University, Bhanjabihar, Ganjam, Odisha, India

Abstract— Haemolymph ascorbic acid of 5th instar larvae of four (4) bivoltine races namely GEN-3, BCON-1, DUN-22, BCON-4 of *Bombyx mori* .L was studied out of four different varieties of bivoltine races GEN-3 race showed best performance in haemolymph ascorbic acid content as well as certain economic characters like cocoon weight (CW), shell weight (SW) and Shell ratio percentage (SR%). Basing upon the above characters the efficiency of the silkworm races was considered. GEN-3 is showing the best performance and highly significant by ANOVA. The correlational study indicates that GEN-3, BCON-1, DUN-22 are highly significant than BCON-4. Their change in concentration of haemolymph ascorbic acid is gradually increased from first day to the 3rd day and it started declining from the 4th day up to the 7th day. The present study showed that there is a direct relationship established between haemolymph ascorbic acid with body weight, cocoon weight and shell weight and shell ratio percentage. The above characters were important for the selection for hybridization as well as commercial rearing.

Key words: Haemolymph Ascorbic Acid, Bivoltine, Cocoon Weight, Shell Weight, Shell Ratio Percentage

I. INTRODUCTION

Haemolymph ascorbic acid is an important component for the metamorphosis of silkworm. The chemical composition of haemolymph is highly variable among the diverse species and at different developmental stages of the some species (Florkin and Jeuniaux, 1974). It constitutes a variety of nitrogenous substances such as amino acids, proteins, amines, ammonia, peptides and ascorbic acid.

One of the most important factors for the proper growth and development of insect species is its nutritional requirements and metabolism in the body. A number of studies have been undertaken to evaluate the vitamin requirements and their metabolism in many lepidopteron insect species.

Besides an attempt has also been made to establish an inter-relationship between morphological, biological and certain economic characters like ascorbic acid content, larval body weight, gland weight and economic characters such as cocoon weight, shell weight and S.R %.

Many researchers showed the larval characters improve by different concentrations of complementary compounds such as ascorbic acid, folic acid, and thiamine. Vitamin-B complex etc. (Sarkar et.al., 1995 ; Nirwani and kaliwal, 1996, 1998; Etabari et.al., 2004).. Several authors also reported these effects about ascorbic acid (Dobzhenok, 1974; Ito, 1978; Singh and Reddy, 1981; EI-Karkasy and Idriss, 1990). Considering the dynamic roles of ascorbic acid in body metabolism an attempt has been made to

evaluate the quantitative changes in ascorbic acid level in seven bivoltine and multivoltine silkworm races of *Bombyx mori.L* Sometimes it is considered as a taxonomic character for the insect (Florkin, M. 1959).

II. RELATED WORK

- 1) It has been well known fact that ascorbic acid is an essential nutrient for many plant feeding insects (Chippendale, G.M. 1975). Akhtar and Asghar (1972), found that vitamins and mineral salts played an important role in the nutrition of silkworm. Ascorbic acid (vitamin C) was studied very early in the history of insect diet.
- 2) Ascorbic acid plays a vital role in participating in various metabolic functions of the body. It has anti-scorbutic and extra- antiscorbutic functions (Hughes, 1977).

Basing on this piece of work many work has been done on the basis of the ascorbic content of silkworm in this context, but my piece of work is a bit different that is a correlational study on the concentration of the haemolymph ascorbic acid of certain bivoltine larvae of the silkworm *Bombyx mori. L* during 5th instar development. It also includes which factors are able to do the concentration change in subsequent varieties of bi and multivoltine also.

III. MATERIAL AND METHODS

Four Bivoltine races of *Bombyx mori.L* namely GEN-3, BCON-1, DUN-22 and BCON-4 races were collected from different breeding stations of Central Silk Board. For the present work silkworms have been collected from Central Sericulture Research and Training Institute (CSRTI), Berhampur (WB). The races were reared and maintained by adopting the Krishnaswami(1978) method. Fifth instar larvae were weighted and sacrificed from the 1st day onwards till cocoon formation. The haemolymph was collected in a clean vile containing few thiourea granules by pricking the abdominal leg just before the last leg. For the estimation of total ascorbic acid content in haemolymph of silkworm the method of Roe (1967) with a little modification was followed. Besides the cocoon weight, the shell weight and shell ratio percentage were calculated. All the data were subjected to ANOVA using standard statistical procedure and the relationships between two continuous variables was computed. The data's were also subjected to correlation coefficient.

IV. OBSERVATION

The Ascorbic acid content in the haemolymph increased significantly to its peak on 3rd day in all the races and it

starts declining from the 4th day up to the last i.e 7th day. (Table 1). Beside haemolymph ascorbic acid progressive significant increments in the body weight were also observed in all the races is a common feature. The present data's showed a significant variation in the quantity of the haemolymph protein was also recorded as in Table-1. The observed data showed strong degrees of positive Correlation between the haemolymph ascorbic acid and the body weight. However a significant correlation was also exists between the cocoon weight and shell weight.

Anova of Bivoltine Races (Haemolymph Ascorbic acid)

Vth Instar Duration	BIVOLTINE RACES			
	GEN-3 MEAN ± S.E.M	BCON-1 MEAN ± S.E.M	DUN-22 MEAN ± S.E.M	BCON-4 MEAN ± S.E.M
1 st Day	0.172 ± 0.003	0.164 ± 0.0002	0.154 ± 0.003	0.148 ± 0.0003
2 nd Day	0.509 ± 0.001	0.491 ± 0.0001	0.368 ± 0.003	0.417 ± 0.0002
3 rd Day	1.002 ± 0.003	0.997 ± 0.003	0.718 ± 0.005	0.621 ± 0.0004
4 th Day	0.8070. ± 0.003	0.705 ± 0.003	0.639 ± 0.0002	0.607 ± 0.0003
5 th Day	0.627 ± 0.004	0.518 ± 0.003	0.541 ± 0.0005	0.535 ± 0.0003
6 th Day	0.513 ± 0.004	0.427 ± 0.006	0.438 ± 0.0024	0.419 ± 0.0003
7 th Day	0.375 ± 0.005	0.366 ± 0.004	0.297 ± 0.0037	0.261 ± 0.0003

NB: NS=Not Significant,
*-- Significant at (0.05) 5% level,
** --Significant at (0.01)1% level

Table 1: Comparison in the Concentration of Ascorbic acid (mg/ml.) in the Haemolymph of some Bivoltine Silkworm Races, Bombyx mori L during 5th instar development

ANALYSIS OF VARIANCE				
Source Of Variation	SS	df	MS	F
Between the Days	1.210921	6	0.20182	45.77955**
Between the Races	0.092735	3	0.030912	7.011763 ^{NS}
Error	0.079353	18	0.004409	
Total	1.383009	27		

Table 2: Coefficient of correlation between Body weight with Haemolymph ascorbic acid of Bivoltine silkworm races Bombyx mori.L

Sl. No	Races of Silkworm	Variables	Coefficient of correlation 'r'	df	't' Test	df
1	GEN-3	BW Vs HASC	0.4303*	06	2.201*	11
2	BCON-1	BW Vs HASC	0.3134*	06	2.201*	11
3	DUN-22	BW Vs HASC	0.4980*	06	2.262*	9
4	BCON-4	BW Vs HASC	0.2409*	06	2.306*	8

BW-Body Weight
GW-Gland Weight
HASC-Haemolymph Ascorbic acid
* Significant at 0.05 level
** Significant at 0.01 level
*** Significant at 0.001 level

V. RESULT AND DISCUSSION

It has been well known fact that ascorbic acid is an essential nutrient for many plant feeding insects (Chippendale, G.M. 1975). Akhtar and Asghar (1972), found that vitamins and mineral salts played an important role in the nutrition of silkworm. Ascorbic acid (vitamin C) was studied very early in the history of insect diet. Many insects are now known to require dietary ascorbic acid.

Many researchers showed the larval characters improve by different concentrations of complementary compounds such as ascorbic acid, folic acid, thiamine. Vitamin-B complex etc.(Sarkar et.al., 1995 ; Nirwani and kaliwal, 1996, 1998; Etabari et.al., 2004).. Several authors also reported these effects about ascorbic acid (Dobzhenok, 1974: Ito, 1978; Singh and Reddy, 1981; El-Karkasy and Idriss, 1990). Considering the dynamic roles of ascorbic acid in body metabolism an attempt has been made to evaluate the quantitative changes in ascorbic acid level in seven bivoltine and multivoltine silkworm races of Bombyx mori.L .Besides an attempt has also been made to establish an inter-relationship between morphological, biological and certain economic characters like ascorbic acid content, larval body weight, gland weight and economic characters such as cocoon weight, shell weight and S.R %




Fig-1.a : Comparison between the amount of Haemolymph ascorbic acid and Body weight of GEN-3 (Bivoltine)




Fig-1.b : Comparison between the amount of Haemolymph ascorbic acid and Single Cocoon weight of GEN-3 (Bivoltine)










Fig-1.c : Comparison between the amount of Haemolymph ascorbic acid and Single Shell weight of GEN-3 (Bivoltine)

Ascorbic acid in the haemolymph of the fifth instar larvae of seven races of Bombyx mori.L was studied. The result had shown a progressive increment in the ascorbic acid content up to the third day of the fifth instar larvae in all the races as in Table-1 and Fig-1 & 2. Thereafter, there was a gradual decline of ascorbic acid content till the end of the fifth instar larvae in all the races is a common phenomena observed as in Table:-1 and Fig-1 & 2. The data were subjected to statistical analysis. There was a significant increment in the ascorbic acid content of the Haemolymph during the feeding phase (1st ~3rd day) of 5th instar larvae of all the races as in Table 1. However, there was a gradual

decrement in the ascorbic acid content from the 4th day onwards.

The data were also subjected to ANOVA test and coefficient of correlation. The result showed a significant variation in the quantity of ascorbic acid among different races of Bombyx mori.L.



Highest recorded value of the ascorbic acid was found in GEN-3 followed by other races among Bivoltine races as in Table-1. However, the amount of ascorbic acid decreased from the fourth days of fifth instar period might be due to the utilization of ascorbic acid for its exponential growth in the synthetic phase.

Variation in the quantity of ascorbic acid are exists among races may be due to their magnitude of consumption and assimilation of mulberry leaves.

Nutrition plays an important role in improving the growth and development of the silkworm, Bombyx mori like other organisms. Legay (1958) stated that silk production is dependent on the larval nutrition which plays a very effective role in producing good quality cocoons. One of the most important micronutrient necessary for the growth and development is ascorbic acid. It plays a vital role in metabolism, participating in various metabolic functions of the body. There is an absolute requirement of ascorbic acid for collagen synthesis. Sengupta et al (1972) showed that the silkworm requires certain essential sugars, proteins, amino acids and vitamins for its normal growth, survival and enhancement of the silk production. It is generally accepted that all insects require vitamins, especially water soluble vitamins, such as ascorbic acid, thiamin, riboflavin, pantothenic acid, biotin, folic acid (Chapman, 1988). Growth retardation caused by lack of each one of these vitamins is rather small, but better growth is obtained by adding these vitamins to their diet (Horie and Ito, 1965; Ito, 1978; Horie, 1995).




Therefore, the necessity of ascorbic acid in the diet of herbivorous insect for feeding requirement and antioxidant activity has been confirmed by many authors (Felton and Summers, 1993 and 1995).

Micronutrients are essential component for the growth and development of silkworm races. Ito (1978) determined that vitamins present in the mulberry leaves satisfy minimum needs of silkworm. The effect of vitamin supplementation on the growth of Bombyx mori have been investigated by many researchers (Majumdar and Medda 1975; Bhattacharyya and Medda, 1981; Das and Medda, 1988; Faruki, 1990, 1998, 2005; Babu et al, 1992; Faruki et al., 1992; Khan and Saha, 1996; Nirwani and Kaliwal, 1996, 1998; Mosallanejad, 2002; Etebari et al, 2004; Rajabi et al., 2006 a,b). Ascorbic acid has many important functions in the animal body. It is a powerful antioxidant, protecting against oxidative damage to DNA, membrane lipid and protein. Antioxidant activity of ascorbic acid decreases reactive oxygen species and oxidative pressure. So, that the absorption of nutritious substances in the midgut would increase (Felton and Summers, 1993). Ascorbic acid has always been regarded as indispensable for the growth and development of Bombyx mori. In fact ascorbic acid is present in large amount in mulberry leaves, the exclusive food for the silkworm, and insects are incapable of synthesizing it.

Ito (1961) recorded the interrelationship between ascorbic acid supplementation and growth of silk worm. Babu et al., (1992) observed that the 1st and 2nd instar larvae reared on 1.5% ascorbic acid enriched mulberry leaves resulted in higher silk filament length, weight and denier values. Chauhan and Singh, 1992 showed that 1% concentration of ascorbic acid could increase in the number of eggs in the silkworm. Many workers established that ascorbic acid enhances the silk yield of the mulberry

silkworm *Bombyx mori*. (A.EI-karaksy and M.I.Idriss, 1990).


The available data's were subjected to coefficient of determination (R²). In (Fig-1.a to 1.l) correlation between Haemolymph ascorbic acid and body weight of fifth instar larvae taken in the analysis. The results were found a low positive correlation, (R²= 0.185) and the regression line obtained $Y = 0.305X + 0.083$. Only 18.5% of correlation were exist between the two variables like ascorbic acid content and body mass of fifth instar period as in Fig-1.a up to 1.l., that indicates only 18.5% of the data is in a interreciprocal influence of both characters. From the data it was evident that if (R² = 0.004) would increased in the ascorbic acid content, it was explained that the body weight also increased in the same direction.



When a correlation was established with ascorbic acid content and cocoon weight and shell weight, which should a very low positive correlation (R² = 0.004) and the regression line obtained $Y = 0.008X + 0.537$ as in Fig-22. Such types of relationship were found in all the races. From the result it was evident that ascorbic acid have a positive effect on the body mass but no such direct effect on the economic parameters like cocoon weight, shell weight.

From the observation it may conclude that GEN-3 race might have better nutrigenetic trait than the BCON-1, DUN-22 and BCON-4 of Bivoltine races (Table-1) basing upon their ascorbic acid content. Presently, nutrigenetic trait analysis is a prime factor while selecting breeds for rearing performances. Therefore, the amount of ascorbic acid present in the larvae might play a crucial role in growth, development, moulting and survivability of mulberry silkworm *Bombyx mori*.L. The amount of ascorbic acid in the fifth instar larval period might have taken as one of the

important criteria for selecting *Bombyx mori*.L. Race for germplasm rearing.



However, considering the quantity of ascorbic acid among *Bombyx mori* races. It may assumed that GEN-3 performed the best. Therefore, the GEN-3 race might have considered to be a better phago-stimulant efficient growth promoter silk worm races. If the mulberry diet is added with ascorbic acid, it is possible to show improvement in sericulture industry. It has already an established fact that ascorbic acid has great role on the growth promoter as well as booster of silk production.(Ito, 1961, Javed and Gondal, 2002; Hussain and Javed., 2002 ; Pravakar,2004 ; Etabari et.al., 2004).

Basing on the amount of ascorbic acid it may be assumed that GEN-3 race is a better nutrigenetic trait. However from the present study explore the possible interrelationship between growth and development with the ascorbic acid content in the haemolymph. Considering many good effects and importance of ascorbic acid as suggested by many authors, it may assumed that amount of ascorbic acid present in the larvae might play a crucial role in growth, development, moulting and survivability of mulberry silk worm *Bombyx mori*.L. Therefore, quantification of ascorbic acid is highly essential for the selection processes.

VI. CONCLUSION

In the present study the amount of the haemolymph Ascorbic acid in fifth instar period from four bivoltine races were recorded to establish a profile of silkworm haemolymph Ascorbic acid. The present data also gave an insight into a comprehensive understanding of silkworm metamorphosis.

ACKNOWLEDGEMENT

The authors thank the authorities of Berhampur University, Odisha, for providing necessary laboratory facilities.

REFERENCES

- [1] Babu, M.; Swamy, M.T.; Rao, P.K. and Rao, M.S.: Effect of ascorbic acid – enriched mulberry leaves on rearing of *Bombyx mori* L. Indian J. Seric. 31(2): 111-114, 1992.
- [2] Chapman, R.F.: The Insect Structure and Function, Cambridge University Press, Cambridge, 1998.
- [3] Chippendale, G.M.: Ascorbic acid an essential for plant feeding insect, *Diatraea grandiosella* Vol.105:4.pp.499-507, 1975.

- [4] El-karakasy, I.R. and Idriss, M.: Ascorbic acid enhances the silk yield of the mulberry silkworm *Bombyx mori* L. *J. Appl. Entomol.* 109: 81-86, 1990.
- [5] Etebari, K. and Matindoost, L.: Effects of hypervitaminosis of vitamin B3 on silkworm biology. *J. Biosci.* 29:417-422, 2004.
- [6] Etebari, K., Ebadi, R. and Matindoost, L.: The effects of vitamin C on biological, biochemical and economical characteristics of the silkworm *Bombyx mori* L. *International J. Indust. Entomol* 8: 81-87, 2004.
- [7] Felton, G.W. and C.B. Summers,: Potential role of ascorbic oxidase as a plant defense protein against insect herbivory. *J. Chem. Ecol.* 19:1553-1568, 1993.
- [8] Felton, G. W. and C. B. Summers: Antioxidant systems in insects. *Arch. Insect Biochem. Physiol.* 29, 187-197, 1995.
- [9] Florin and Jeuniaux: Haemolymph composition. In the *Physiology of Insect* (Ed by Rockstein, M) Academic Press, New York, 5: pp. 255-307, 1974.
- [10] Florin, M. & Jeuniaux, C. :*The physiology of Insecta.* V. Ed. Rockstein Academic Press, 1974.
- [11] Horie, Y; Watanabe, K. and Shinohara, E.: Effects of diets on body weight, silk gland weight and blood composition in the silkworm *Sansi Kenkyu* (*Acta Sericologia*), 78: 44-5, 1971.
- [12] Ito, T.: Effect of dietary ascorbic acid on the silkworm *Bombyx mori* L. *Nature*, 192: 951-952, 1961.
- [13] Ito, T. (1978) : *Silkworm nutrition; In the Silkworm an Important Laboratory Tool.* Tazima, Y. (Ed.), pp. 121-157, Kodansha Ltd, Tokyo ,1978..
- [14] Ito, T. and Arai : Nutrition of silkworm *Bombyx mori* L. IX. Further studies on the nutritive effects of ascorbic acid. *Bull.Seric. Expt. Statn.*, 20: 1-19, 1965.
- [15] Ito, T. and Horie, Y.,: Carbohydrate metabolism of the midgut of the silkworm, *Bombyx mori* L. *Archives of biochemistry & biophysics*, 80, pp 174-176, 1959.
- [16] Ito, T. and Mukayama, F.: Relationship between protein content of artificial diet and cocoon quality in the silkworm. *Sansi Kenkyu* (*Acta Sericologia*), 77. pp. 76-81, 1970.
- [17] Ito, T and Tanaka, M.: Nutrition of the silkworm, *Bombyx mori* L.VI. Effect of Concentration of sugar and protein added in artificial diet. *Seri.Exp.Stn.Japan.*18: 1-34, 1962.
- [18] Legay, J.M.: Recent advances in silkworm nutrition. *Ann. Rev. Ent.*, 3:75- 124, 1958.
- [19] Majumdar, A.C. and Medda, A.K.: Studies on the thyroxine and vitamin B12 induced changes in the life cycles of silkworms. *Indian J. Physiol. Appl. Sci.* 29:199-203, 1975.
- [20] Sarkar, A.; Haque, M.; Rab. M. and Absar, N.: Effect of feeding mulberry (*Morus sp.*) leaves supplemented with different nutrients to silkworm, (*Bombyx mori* L.). *Curr. Sci.* 69(2): 185-188, 1995.
- [21] Suresh Kumar N, Yamamoto T, Basavaraja HK, Datta RK: Studies on the effect of high temperature on F1 hybrids between Polyvoltine and bivoltine silkworm races of *Bombyx mori* L. *International Journal of Industrial Entomology*2, 123-127, 2001.
- [22] Shigematsu, H.: Synthesis of blood proteins by the fat body in the silkworm, *Bombyx mori* L. *Nature*, 182:880-882, 1958.