

Adverse Impact of Air Pollutants on Human Health: A Case Study of Aurangabad City

Dr. Vijaykumar Bhikusing Pawar¹ Dr. Kshama Khobragade²

¹Research Scholar ²Reader & Head of Department

²Department of Environmental Science

^{1,2}S. B. E. S. College of Science, Aurangabad. 431 001, India

Abstract— Exposure to air pollution is an inescapable part of our urban life. In this study, the interaction patterns of air pollutants, SO₂, NO_x, SPM and PM₁₀ are investigated based on measured database of the study area in Aurangabad. The paper mainly concentrates on the Impacts of Indoor and outdoor air pollutants especially on health, with result that Human Hygienic may be deteriorated, economic productivity may be declined. It has been concluded that intervention of the government has become indispensable to implement strong environmental policy and taking care of growing threats on air pollution in more institutionalized, regular and automatic basis is cannot be avoided.

Key words: Airborne Pollutants, Health Impact, Industrial Air Pollution, Air Quality Assessment

I. INTRODUCTION

Aurangabad is one of the fastest developing city in India. The location co-ordinates for Aurangabad are N 19° 53' 47" - E 75° 23' 54" and located 512 meters above sea level. The city is surrounded by hills on all sides.

Air pollution has adverse impact on human health as well as the health of other living entities and stress vegetation. Depending upon the life time of the pollutants, location of the sources and prevailing air currents, receptors, may be located at time intervals from near instantaneous to several decades¹ good air quality is essential for health of people and the environment, although the significant improvement have been made in many countries over the last 2-3 decades, air quality particularly in urban areas, remain a priority issue on most national environmental agendas² particulates and gaseous emission of pollutants from industries and auto-exhaust is responsible for rising discomfort, increasing airway diseases, decreasing productivity and deterioration of artistic and cultural patrimony in urban centers³ The presence of air pollutants over the prescribed limit in the lower atmosphere is not only injurious to humans but also to animals, foliage, fruits, vegetables and microbial life and may even damage property⁴ urban air pollution is a major focus of public health concern and regulatory activity⁵

The purpose of this study is to

- a) Quantify air quality,
- b) Observe health effects.

II. MATERIALS & METHODS

The study is carried out

- 1) To study concentration of few air pollutants
- 2) To estimate the adverse health effects due to air pollution.
- 3) To study relation of air pollutants with that of various diseases occurred in different age groups of both genders.

- 4) To quantify the cases of premature death and morbidity associated with air pollution through the survey questionnaire collected in Aurangabad.

Annual temperature in Aurangabad ranges from 9° to 40° C. Average annual rainfall is 725 mm. Various Industries like Chemicals, Pharmaceuticals Drug, Distilleries, oil refineries, fertilizer plants, Pulp and Paper etc. are situated in this area. This study covers pollutants viz. SO₂, NO_x, SPM and PM₁₀ and these are investigated based on the measured database of Maharashtra Pollution Control Board (MPCB).

III. CROSS-SECTIONAL SURVEY

A survey questionnaire relating to air pollution, health effects, local economic status and other factors and suggestions for improvement of the environmental status, was circulated for public perception and to gain a better understanding of public opinion. The survey was based on a mix of written responses to the questionnaire as well as personal interviews. The survey was conducted in 13 locations during June to December 2016. Hundred respondents, both adults and children living in Aurangabad and the surrounding area were surveyed.

A. Air Pollution

Biofuels is an issue that requires to be addressed through health policy. Some of the highest concentrations of pollutants occur due to the use of bio-fuels like Wood, Agriculture crop residues and dung cake for cooking in rural environment, Automobile industries, drug and distilleries are main sources of air pollution and their emission break out various air borne diseases like bronchitis, asthma, pneumonia and even lung cancer and skin diseases and Vehicular Emission has been identified as the greatest environmental danger in a number of cities in developing countries as well as Burning of agricultural harvest waste causes problems related with air pollution like Smoke nuisance ,Effects on human hygiene, Threat to wild life, Damage to historical monuments.

The pollutants which are emitted by selective industries like sugar factories, distilleries, Cement factories, paints, pulp and paper industries have damaging effects on human health, Forest, Cattle and Bio-diversity.

IV. RESULTS & DISCUSSION

A. Morbidity Data – Findings

Morbidity data were collected from major hospitals of Aurangabad, which consists of bronchial asthma, respiratory disorders, eye conjunctivitis, scabies and other skin infections.




Fig. 1: Bronchial Asthma Cases from 2016 to 2017

(Source: As per major five hospital records in Aurangabad city of year 2016- 2017)

B. Asthma Cases

Season wise hospital admissions for treatment of asthma are shown in Table No.(1) from April 2016 to December 2017 in the study, Generally, both adult male (am) and adult female (af) were affected by asthma. Females are more affected. Though both male and female children (mc and fc) were

Year	Season	Adult Male (am)	Adult Female (af)	Male Child (mc)	Female Child (fc)	Month with high rate of Admission
2016	Summer	122	201	162	134	June
	Monsoon	133	354	177	156	Aug.
	Winter	139	365	186	163	Feb.
2017	Summer	314	296	175	326	May
	Monsoon	301	294	183	324	Aug
	Winter	299	279	191	321	Nov.

Table 1: Hospital Admissions for Treatment of Asthma Cases – Season-Wise Maximum

(Source: As per five Major Hospital Records in Aurangabad city for year 2016 and 2017)

Year	Season	Adult Male	Adult Female	Male Child	Female Child	Month With High Rate of Admission
2016	Summer	53	74	Na	Na	Na
	Monsoon	302	710	37	93	Jul. to Sept.
	Winter	39	136	488	360	Very High During All Months
2017	Summer	318	469	187	255	During All Months
	Monsoon	414	138	325	474	Very High During All Months
	Winter	45	262	35	241	Nov. to Feb.

Table 2: Hospital Admissions for Treatment of Respiratory Cases – Season Wise Maximum

NA - Not Available

(Source: As per major five hospital records in Aurangabad city of year 2016-2017)

D. Eye & Skin Problem Cases

In November 2016, both adults and children were affected with eye problems while in Feb. 2017, female and male children were more affected with skin problems.

E. Mortality Data – Findings

Data regarding death were collected from the Town Aurangabad., based on birth and death records from 2016 to June 2017. Table 3 shows the year and gender-wise death cases that occurred in Aurangabad households under various age groups. It can be noticed that the total number of deaths recorded have been consistently on an increase each year. It is also generally observed that the number of death cases that occurred among males is higher than females. A comparison of the number of death cases with that of the age groups

affected, hospital admissions with respect to female children are higher.

In May 2016, the maximum value of PM₁₀ pollutant was 219 $\mu\text{g}/\text{m}^3$ and that of SO₂ was 26 $\mu\text{g}/\text{m}^3$ according to the air quality data. Comparison of air quality data with morbidity data reveals that hospital admissions for treatment of asthma were as high as am 55; af 62; mc 40; fc 50. It was further observed that the total number of cases recorded for treatment of various diseases for new cases were 2998, old cases 1554, as against the total population of the Aurangabad city.

C. Respiratory Problem Cases

Season wise distribution of hospital admissions for treatment of respiratory problems is given in Table 2. Generally, both adult male and female were affected by respiratory problems. Females are more affected and were considerably in higher numbers for treatment. As far the children, both male and female children were affected in winter 2016⁶. The maximum value of PM₁₀ pollutant was 276 $\mu\text{g}/\text{m}^3$, NO_x 81 $\mu\text{g}/\text{m}^3$, SO₂ 130 $\mu\text{g}/\text{m}^3$ According to the air quality data it was further observed that the air quality data it was further observed that the total number of cases recorded for treatment of various diseases for new cases were 3280, old cases 1868, as against the total population of the area

indicated that the number of death cases in 40–59 age group of both male and female category are high, followed by the age group 70. It is seen that ten male and six female death cases occurred particularly in May 2016.

It is further observed that the percentage of deaths due to heart attacks is considerably high, compared with other causes for death as shown in Table No. 3 The air pollutant PM₁₀ is of prime concern, as it triggers increase in cases with breathing problems, asthma and heart attacks.²

F. Cross-Sectional Survey

Among the villages that intersect Aurangabad. The inhabitants of Cidco, Chikalthana and Waluj-Pandharpur areas were severely affected by asthma, respiratory problems were noticed at the rate of 8 to 10 persons in every studied

area. Similarly, 4 to 8 persons were affected by the same problem in Chikalthana, In the remaining areas, an average of 2 to 3 persons were affected while conducting the survey in December 2016, the respondents informed that 2 to 3 cases of premature deaths had occurred in Cidco, Chikalthana, Waluj-

Pandharpur and Shendra The survey result indicates that in the said villages, the urban poor people spent at the rate of Rs. 150 to 200 per month per head towards cost of illness, while others spent at Rs 250 to 300.

		Death cases in numbers						
		Age Group						
Year	Gender	0-9	10-19	20-39	40-59	60-69	>70	Total
2016	Male	1	0	8	20	14	19	62
	Female	2	3	3	3	4	12	27
2017	Male	1	1	4	20	12	14	52
	Female	0	1	4	8	4	18	35

(Source: As per Major Hospital Records in Aurangabad city of year 2016-2017)

Table 3: Death Cases that Occurred in Aurangabad Households from 2016 to June 2017

V. CONCLUSION

This is a study about higher air pollutant concentrations, which give rise to significant adverse health effects, and it is an issue of serious public health threat to inhabitants of the study area. However, increasing levels of pollutants, due to industrial activities and alarming increase in the number of vehicles may become a problem in the future, with greater health impact. A comparison of the survey results of morbidity and mortality cases indicates that the impact of SPM and PM₁₀ levels gives rise to increased hospitalization for respiratory problems like asthma and gradual increase in mortality cases. Cases of premature death due to heart attack are on the increase in males under the age group 40–59 compared to all other causes for mortality. The study concludes that the seriousness of the air pollution problem in Aurangabad. May aggravate further, if not brought under control in this scenario, the city development and environmental enforcement authorities have to adopt future strategies to combat the menace of air pollution.

VI. SUGGESTIONS

- a) Indoor air pollution control strategies
 - 1) Improving the ventilation in the cooking area so as to reduce particulate matter concentration
 - 2) Increasing access to clean fuels such as biogas, solar stoves, liquid petroleum gas etc.
- b) Outdoor air pollution control strategies
 - 1) Manufacturer of such Vehicles which do not pollute the atmosphere beyond permissible limits
 - 2) Serious actions and laws over burning Agriculture waste
 - 3) Sources of Industrial pollution must be taken into account in setting Air Quality standards
 - 4) Producers and consumers should be aware about the air pollutants and their hazardous impacts

REFERENCES

[1] Chattopadhyay B. P. et. al., Respiratory Health status of the road side school children at Kolkatta, Journal of environmental Science and Engineering, Vol. 47 No. 3,(July 2005) pp.202-211

[2] Evans F. and Smith K., Do new health conditions support mortality air pollution effects? J. Environ. Econ. Manage. [on line from <http://www.sciencedirect.com>]

[3] Matani A. G., Vehicular Pollution Poisoning the air in cities, Ecology, Environment and Conservation, Vol. no. 4, (1998) pp. 295-297

[4] MPCB, Ambient air quality status (2016–2017). National ambient air quality monitoring series: NAAQMS/2016-2017, Maharashtra Pollution Control Board, Mumbai. [on line from <http://www.mpcb.com>]

[5] National Environment Policy, Draft for Comments: 30 November 2004.

[6] Pandey, D. K., Effects of air pollution on health. Employment News, 2001, vol. XXVI, pp. 17–23.

[7] Rao M. N. and Rao H. V. N., In Air Pollution, Tata McGraw Hill, New Delhi, (1997) p. 43.